

The response of the X-ray luminosity to the interaction strength in Seyfert 1 & 2 galaxies

F. Pfefferkorn & T. Boller^a

^a*Max-Planck-Institut für extraterrestrische Physik, Giessenbachstrasse, 85748 Garching, Germany email: pfefferk@mpe.mpg.de & bol@mpe.mpg.de*

Abstract

We have correlated a sample of 99 Seyfert 1 & 98 Seyfert 2 (Rafanelli et al., 1995) interacting galaxies with the survey and pointed ROSAT data and have detected 91 Seyferts 1 & 47 Seyfert 2s in the X-ray band. We have performed spectral and timing analysis of the X-ray detected sources, and we have examined the relation of the X-ray luminosity L_X with the interaction strength Q .

Key words: galaxies: active, Seyfert 1, Seyfert 2, merger; X-rays: galaxies

1 Introduction

Interacting galaxies are considered to have a wide importance on triggering of starburst and AGN activity. Therefore, we have investigated two subsamples of spectroscopically-selected, X-ray detected interacting Seyfert 1s, including NLS1, and Seyfert 2s galaxies (Rafanelli et al., 1995) with $z < 0.11$, $m \leq 15.5$ and $\delta \geq -23^\circ$.

The pairs have been selected on POSS plates using the following criteria given by Rafanelli et al. (1995): separation between the components $S < 3D_p$, where D_p is the apparent major axis of the Seyfert galaxy, and a magnitude-difference between Seyfert and the companion $\Delta m = m_{comp} - m_{seymert} < 3$.

The identification is based on X-ray contour lines overlaid on optical images taken from DSS. In Fig. 1 we show as an example the overlay from the NLS1 galaxy Mkn 896. Fig. 2 shows the spectrum of this object. The deviation from the fit at energy 0.7 keV in Fig. 2 indicates the presence of a warm absorber. The pointing and survey lightcurves in Fig. 3 show the variability of this source on short time scales.




Fig. 1. The NLS1 galaxy Mkn 896 with a companion galaxy. The X-ray contours are overlaid on the optical image.




Fig. 2. Simple power-law fit results to the NLS1 Mrk 896.

2 First results on the interaction strength Q

To derive the interaction strength Q , we concentrate on the tidal force per unit mass produced by a companion on a primary galaxy, which is proportional to $M_c \cdot R^{-3}$. M_c is the mass of the companion and R is its distance from the center of the primary galaxy. In most cases, M_c and the absolute value of R




Fig. 3. The lightcurves from the pointing and survey observation shows the short-time variability of Mkn896.

are unknown. Instead, these parameters are related to the dimensions of the pair. Rubin et al. (1982) describe the dependence of the mass M of a galaxy on the size of its major axis d as $M \propto d^\gamma$ and we use $\gamma = 1.5$ (Dahari, 1984). If we utilize the apparent diameter of the primary galaxy D_p as a scaling factor, we obtain:

$$M_c \propto (D_c/D_p)^{1.5} \quad \text{and} \quad R \propto S/D_p$$

Using these relations, we get as the dimensionless gravitational interaction strength Q :

$$\frac{M_c}{R^3} \propto \frac{(D_c \cdot D_p)^{1.5}}{S^3} \equiv Q$$

This parameter is obviously large for close and relatively large companions.

Fig.4 shows interaction strength Q vs. the X-ray luminosity for Seyfert 1 galaxies. Noticeable is a faint tendency for the luminosity increase for increasing values of interaction strength. A possible explanation for the low luminosity sources are the presence of a high hydrogen column density in the nucleus and the emission from the circumnuclear starbursts.

The galaxy interactions or mergers are able to trigger a burst of star formation, which can be a reason for the rise in luminosity. Several studies about physical mechanisms which could trigger a burst of strong star formation have been investigated by Barnes & Hernquist (1991), Jog & Das (1992) and Jog & Solomon (1992). Also many observations suggest that galactic encounters enhance star formation rates, as demonstrated by studies of the optical (e.g., Kennicutt et al. 1987; Bushouse 1987), infrared (e.g., Bernlöhner 1993; Telesco et al. 1993) and radio (e.g., Hummel 1981; Smith & Kassim 1993) emission from interacting systems.

For each source detected in the All-Sky Survey with more than 100 counts in an exposure time of 400 sec and for all sources within pointed observations, we




Fig. 4. Soft X-ray luminosity L vs. dimensionless gravitational interaction strength Q for the Seyfert 1 galaxies. The plot suggests an increase in luminosity for increasing values of the interaction strength. The luminosity has been calculated from the absorption corrected flux, integrated over the energy range $0.1 - 2.4 \text{ keV}$ and using the Hubble constant $H_o = 75 \frac{\text{km}}{\text{s Mpc}}$.

performed a spectral analysis and created a lightcurve. Studies of variability were possible, if more than one pointed observation was available or if both a detection in the survey and in a pointed observation existed. We are extending our studies on starburst galaxies to examine the influence of interaction with companions on the X-ray luminosity. A paper with data concerning X-ray properties, luminosity, flux, count rate variation and lightcurves for Seyfert 1s & 2s is in preparation.

References

- [1] Barnes J.E., & Hernquist L.E., *Astroph. J.* **370** (1991) L65
- [2] Bernlöhr K., *A&A* **268** (1993) 25
- [3] Bushouse H.A., *Astroph. J.* **320** (1987) 49
- [4] Dahari O., *Astron. J.* **89** (1984) 966
- [5] Hummel E., *A&A* **96** (1981) 111
- [6] Jog C.J., & Das M., *Astroph. J.* **400** (1992) 476
- [7] Jog C.J., & Solomon P.M., *Astroph. J.* **387** (1992) 152
- [8] Kennicutt Jr. R.C., Keel W.C., van der Hulst J.M., Hummel E., & Roettiger K.A. *Astron. J.* **93** (1987) 1011
- [9] Rafanelli P., Violato M., & Baruffolo A., *Astron. J.* **109** (1995) 1546
- [10] Rubin V.C., Ford Jr. W.K., Thonnard N., & Burstein D., *Astroph. J.* **261** (1982) 439
- [11] Smith E.P., & Kassim N.E., *Astron. J.* **105** (1993) 46
- [12] Telesco C.M., Dressel L.L., & Wolstencroft R.D., *Astroph. J.* **414** (1993) 120